

INTEGRATED YOGA AS A FIRST-LINE TREATMENT FOR ANTENATAL AND POSTNATAL MENTAL HEALTH

Dr Mahi Mishra (MS O&G)¹ Dr Tanmay Bagade²

¹Reader (Department of O&G), Government Autonomous Ayurveda College and Hospital, F-1, Govt. Ayurveda College Campus, Near Mandre Mata Mandir, Kampoo, Lashkar Gwalior (M.P.) 474009, India, ²Lecturer (Medical education, Global Health) at The University of Newcastle (UON), University Drive, Callaghan, NSW 2308, Australia. Office: Level 4 West(PublicHealth), HMRI Building, John Hunter Hospital Campus

DOI: <https://doi.org/10.47071/pijar.2022.v07i05.01>

ABSTRACT:

Postnatal period of women is one of the most stressful events of a woman's life. The journey from a pre-pregnant state to day to day challenges of managing a new born baby is not only stressful but also leads to several mental health issues in women during pregnancy and postnatally. Major Depressive Disorders (MDD) are becoming a global burden of diseases amongst women of child-bearing age. 1 "Major depressive disorder (MDD) is a mental disorder characterised by at least two weeks of pervasive depressed mood, loss of pleasure in daily activities, weight loss, insomnia, agitation, fatigue, feelings of worthlessness or guilt, attentional problems, thoughts of death, and suicidal ideation".

Key words: Yoga, Pranayama, Meditation Post-Natal Depression (PND), Major Depressive Disorders (MDD), Maternal mental health, Cognitive Behavioural Therapy (CBT), Interpersonal Psychotherapy (IPT),

PARYESHANA

INTRODUCTION:

In 2017, the MDD was found to be prevalent in 11.9% women during the

perinatal period in a systematic review.

The prevalence of Post-natal

Depression (PND) is also estimated to be more than 15%, which is much higher than MDD during the antenatal period. In developing countries, an estimated 15.6% of women suffer from antenatal depression and 19.8% suffer from PND. WHO estimates that 20% of mothers in developing countries suffer from clinical depression. MDDs adversely affect women's well-being as well as their children's growth and development. In low-income countries of South Asia, poor maternal mental health is associated with decreased infant growth. Maternal mental health disorders are treatable and have become an international healthcare priority. However, there is a wide range of choices available to the patients seeking care for PND.

Yoga description review

Currently, numerous clinical and non-clinical treatments are available to manage MDDs. The most common treatments are Cognitive Behavioral Therapy (CBT), Interpersonal Psychotherapy (IPT), talk therapies, collaborative care models, complementary and alternative medicine approaches, light therapy, brain stimulation, and

psychopharmacological interventions. CBT and IPT have shown promising results in managing MDDs. However, due to ethical issues in conducting human trials on pregnant mothers, the effectiveness of most of these interventions is studied using small sample sizes; therefore the generalisability of these studies is questionable. Pharmacological interventions are mostly studied on animal models. Gaps in the literature on the efficacy of clinical and non-clinical interventions for PND have posed a significant challenge for the clinicians as well. Some studies have encouraged a tailored approach to treat PND. A discrete choice experiment in postnatal women in Australia revealed that women who had higher education chose effective treatment. Cost and treatment type was another critical predictor for women to choose a particular treatment. In this study, Yoga and meditation was one of the most popular and preferred choices of treatment for postnatal women suffering from depression. Yoga is one of the most cost-effective exercises used in the Indian sub-continent for thousands of years. Yoga, which includes yogic postures, pranayama

(breathing exercises), and meditation, is a simple, effective, and cost- a form of intervention that can be used to treat antenatal, as well as post-natal depression. Yoga has gained popularity globally since the last decade and has initiated numerous clinical studies across the world. Internationally, yoga- related interventions are grouped into two types: physical-exercise-based yoga and integrated yoga which, besides physical exercises, includes pranayama, meditation or deep relaxation. Physical-exercise-based yoga is an incomplete type of yoga and should not be recommended for managing any mental health-related disorders. Gong et al., in systematic review and meta-analysis of six randomized controlled trials (RCTs) have concluded that 'Integrated Yoga' significantly reduces post-natal depression. In another review about health effects of yoga on pregnancy, Jiang et al. revealed that ten RCTs indicated yoga showed significant reduction in incidence of prenatal mental health disorders, lower levels of pain and stress, lower rate of small for gestational age child, and higher score of relationship.

Result

Yoga can be safely recommended for antenatal and postnatal lumbopelvic pain. Yoga has also shown a significant reduction in cortisol levels and other stress markers in other studies involving women with antenatal and postnatal depressive disorders.

Discussion

Yoga benefits physical and mental health via down-regulation of the hypothalamic-pituitary- adrenal (HPA) axis and the sympathetic nervous system (SNS). Down regulation of HPA axis can lead to the reduction of non-communicable diseases such as diabetes, cardiovascular diseases, obesity, and depression. A literature review of Yoga with other forms of exercise elaborated on the effectiveness of Yoga in comparison to other forms of exercises. In healthy, as well as diseased population, Yoga was found to be significantly effective in improving several health outcomes such as the reduction in blood glucose and lipid levels, salivary cortisol levels, and oxidative stress. Yoga also had a positive effect on reducing sleep disorders.

Conclusion

Yoga is also more effective than walking or other standard prenatal exercises. In comparison to walking, Yoga is significantly effective in improving mood and alleviating anxiety. Cognitive behavioural therapy (CBT) aids patients to modify irrational beliefs and specific cognitive distortions, whereas, Yoga focuses on a holistic approach that combines spiritual, physical, and mental health, along with dealing with private events through meditation. Some studies have shown significantly enhanced and cumulative effect if Yoga is used in combination with CBT in stress management and treating depressive disorders.

Summary

This short review is therefore in favour of recommending Integrated Yoga (yogic postures as physical exercise, pranayama, as breathing exercise, and meditation) for antenatal and postnatal depressive disorders routinely as the first line of treatment. Further comparative studies are needed to see the effect of other forms of therapies with or without the combination of Yoga.

Future Agenda

Longitudinal studies are also needed to analyse the effect of Yoga during the perinatal period on the mental health of the child.

Funding

This study received no external funding. Declaration of interests We declare no competing interests.

References

1. Woody CA, Ferrari AJ, Siskind DJ, Whiteford HA, Harris MG. A systematic review and meta- regression of the prevalence and incidence of perinatal depression. *Journal of Affective Disorders* 2017; 219: 86-92.
2. Kandel H, Pfaff DW. Chapter 1 - Arousal. In: Fink G, ed. *Stress: Physiology, Biochemistry, and Pathology*: Academic Press; 2019: 1-18.
3. Almond P. Postnatal depression: A global public health perspective. *Perspectives in Public Health* 2009; 129(5): 221-7.
4. WHO. Maternal mental health. 2020. https://www.who.int/mental_health/maternal-child/maternal_mental_health/en/ (accessed 18 Jan 2020).

5. Patel V, Rahman A, Jacob KS, Hughes M. Effect of maternal mental health on infant growth in low income countries: new evidence from South Asia. *BMJ* 2004; 328(7443): 820-3.
6. Ride J, Lancsar E. Women's preferences for treatment of perinatal depression and anxiety: A discrete choice experiment. *PLoS ONE* 2016; 11(6).
7. Nillni YI, Mehrizade A, Mayer L, Milanovic S. Treatment of depression, anxiety, and trauma-related disorders during the perinatal period: A systematic review. *Clinical Psychology Review* 2018; 66: 136-48.
8. Vogel L. Tailored treatment for postpartum depression. *Canadian Medical Association Journal* 2011; 183(16): E1163-E4.
9. Gong H, Ni C, Shen X, Wu T, Jiang C. Yoga for prenatal depression: a systematic review and meta-analysis. *BMC Psychiatry* 2015; 15(1): 14.
10. Jiang Q, Wu Z, Zhou L, Dunlop J, Chen P. Effects of Yoga Intervention during Pregnancy: A Review for Current Status. *American Journal of Perinatology* 2015; 32(6): 503-14.
11. Field T, Diego M, Delgado J, Medina L. Yoga and social support reduce prenatal depression, anxiety and cortisol. *J Bodywork Mov Ther* 2013; 17(4): 397-403.
12. Bershadsky S, Trumpfheller L, Kimble HB, et al. The effect of prenatal Hatha yoga on affect, cortisol and depressive symptoms. *Complementary Therapies in Clinical Practice* 2014; 20(2): 106-13.
13. Ross A, Thomas S. The health benefits of yoga and exercise: a review of comparison studies. *The journal of alternative and complementary medicine* 2010; 16(1): 3-12.
14. Carroll JE, Teti DM, Hall MH, Christian LM. Maternal Sleep in Pregnancy and Postpartum Part II: Biomechanisms and Intervention Strategies. *Current Psychiatry Reports* 2019; 21(3).
15. Chris C. Streeter, Theodore H. Whitfield, Liz Owen, et al. Effects of Yoga Versus Walking on Mood, Anxiety, and Brain GABA Levels: A Randomized Controlled MRS Study. *The Journal of Alternative and Complementary Medicine* 2010; 16(11): 1145-52.
16. Vorkapic CF, Rangé B. Reducing the Symptomatology of Panic Disorder: The Effects of a Yoga Program Alone

and in Combination with Cognitive-Behavioral Therapy. *Frontiers in Psychiatry* 2014; 5(177).

17. Granath J, Ingvarsson S, von Thiele U, Lundberg U. Stress Management: A Randomized Study of Cognitive Behavioural Therapy and Yoga. *Cognitive Behaviour Therapy* 2006; 35(1): 3-10.

18. Khalsa MK, Greiner-Ferris JM, Hofmann SG, Khalsa SBS. Yoga-Enhanced Cognitive Behavioural Therapy (Y-CBT) for Anxiety Management: A Pilot Study. *Clinical Psychology & Psychotherapy* 2015; 22(4): 364-71.

Corresponding author:

Dr Mahi Mishra

Reader (Department of O&G), Government Autonomous Ayurveda College and Hospital, F-1, Govt. Ayurveda College Campus, Near Mandre Mata Mandir, Kampoo, Lashkar Gwalior (M.P.) 474009, India

drmahimishra@gmail.com

Source of Support: NIL
Conflict of Interest : None declared

Published BY:
Shri Prasanna Vitthala Education and Charitable Trust (Rea)